Ultra Benfotiamine

Healthy Peripheral Circulation and AGE inhibition†

DESCRIPTION

Ultra Benfotiamine contains the fat-soluble thiamine vitamin called benfotiamine, plus activated vitamin B6, alpha lipoic acid, grape seed extract and pine bark extract to support healthy peripheral circulation and the nervous system, normal kidney function, and AGE (advanced glycation end-product) inhibition that is related to healthy blood sugar metabolism.†

FUNCTIONS

BenfoPure® is a unique proprietary form of Benfotiamine, a fat-soluble form of Vitamin B-1 Thiamine. Studies show that BenfoPure® Benfotiamine is significantly more bioavailable than traditional forms of Vitamin B-1 or thiamine. In Europe, benfotiamine has been utilized for decades in reducing the negative impact of sustained high blood sugar levels.† Benfotiamine supports the body through blocking biochemical pathways by which high blood sugar damages cells. Excessive sugar intake and high blood sugar are associated with a wide range of health complications.

Studies have demonstrated the following benefits:

- Benfotiamine helps support normal peripheral circulation†
- Benfotiamine may reduce discomfort and dysfunction, helping to maintain a healthy nervous system while protecting against oxidative stress in the brain.†
- Benfotiamine supports normal kidney function.†
- Benfotiamine maintains normal endothelial function by reducing the risk of AGEs, which is associated with optimal cardiovascular health.†

Advanced glycation end products (AGEs) are proteins or lipids that become glycated after exposure to sugars. The presence and accumulation of AGEs in many different cell types affect extracellular and intracellular structure and function. In high blood sugar environments and in natural aging, AGEs alter cell structure and normal vascular function.

Vitamin B6 has been shown to help provide comfort for peripheral nerves, especially in those individuals with pyridoxine deficiency. Pyridoxal-5-Phosphate (P5P) is the active form of pyridoxine that the body utilizes directly without need for conversion from pyridoxine. Alpha lipoic acid (ALA) is among the well-researched nutrients for support of peripheral nerve function. ALA may normalize nitric oxide levels to support increased circulation to the neurons. One study confirmed ALA resulted in significant reductions in lipid peroxidation. Proanthocyanidins are a class of potent antioxidants found in pine bark extract, grape seed extract, and many other plants. OPC (oligomeric proanthocyanidins) has been shown to neutralize free radicals, support the strength and structure of blood vessels, and support the normal inflammatory response.† MegaNatural® BP Grape seed extract is different from other grape seed extracts and made by a patented water extraction process that yields a high grade extract with a low molecular weight. Mega Natural BP extract contains over 90% proanthocyanidins. Studies in humans also show it can help maintain normal arterial function and endothelium cell function.†

INDICATIONS

Ultra Benfotiamine is indicated for individuals that desire healthy outcomes of blood sugar metabolism such as healthy nerve function, kidney function, and vascular health.†

FORMULA (#201815-60X)

Ultra Benfotiamine

Healthy Peripheral Circulation and AGE inhibition†

(as BenfoPure®)

Alpha Lipoic Acid 100 mg

MegaNatural®-BP

Grape Seed Extract 25 mg

(Vitis vinifera) (standardized to 90% polyphenols)

Enzogenol® Pine Bark Extract .. 10 mg

Other Ingredients: Hydroxypropyl methylcellulose (capsule), cellulose, dicalcium phosphate, vegetable stearate, silica.

SUGGESTED USE

Adults take one capsule twice daily or as directed by a healthcare professional.

SIDE EFFECTS

No adverse effects have been reported.

STORAGE

Store in a cool, dry place, away from direct light. Keep out of reach of children.

REFERENCES

Nikolić A, Kacar A, Lavrnić D, Basta I, Apostolski S. Srp Arh Celok Lek. 2009 Nov-Dec;137(11-12):594-600. Serbian. PubMed PMID: 20069914. [Benfotiamine].

Haupt E, Ledermann H, Köpcke W. Int J Clin Pharmacol Ther. 2005 Feb;43(2):71-7. PubMed PMID: 15726875. [Benfotiamine].

Du X, Edelstein D, Brownlee M. Diabetologia. 2008 Oct;51(10):1930-2. PubMed PMID:18663426. [Benfotiamine].

Schupp N, Dette EM, Schmid U, Bahner U, Winkler M, Heidland A, Stopper H. Naunyn Schmiedebergs Arch Pharmacol. 2008 Sep;378(3):283-91 PubMed PMID: 18509620. [Benfotiamine].

Schmid U, Stopper H, Heidland A, Schupp N. Diabetes Metab Res Rev. 2008 Jul-Aug;24(5):371-7. PubMed PMID: 18384109. [Benfotiamine].

Stirban A, et al. Diabetes Care. 2006 Sep;29(9):2064-71. PubMed PMID: 16936154. [Benfotiamine].

Alternative Medicine Review Volume 11, Number 4 December 2006.

Nagamatsu M, Nickander KK, Schmelzer JD, et al. Diabetes Care 1995;18:1160-1167. [Lipoic acid].

Abbas ZG, Swai AB. East Afr Med J 1997;74:803-808. [Thiamine and Pyridoxine].

Yokozawa T, Cho EJ, Park CH, Kim JH. Evid Based Complement Alternat Med. 2012;2012:623879. Epub 2011 Sep 7. [Proanthocyanidins].

Edirisinghe I, Burton-Freeman B, Tissa Kappagoda C. Clin Sci (Lond). 2008 Feb;114(4):331-7.

Shand B, et al. Phytother Res. 2003 May;17(5):490-4. [Enzogenol].

Ultra Benfotiamine

Healthy Peripheral Circulation and AGE inhibition†

Walker MJ Jr, Morris LM, Cheng D. Rev Neurol Dis. 2010;7(4):132-9. [L-methylfolate, methylcobalamin, and pyridoxal 5'-phosphate].

Ying Z, Xie X, Chen M, Yi K, Rajagopalan S. Vascul Pharmacol. 2014 Nov 22. pii: S1537-1891(14)00192-X. doi: 10.1016/j.vph.2014.11.004. [Alpha lipoic acid].

Ferreira PM, Militão GC, Freitas RM. Neurosci Lett. 2009 Oct 23;464(2):131-4. doi:10.1016/j.neulet.2009.08.051. [Lipoic acid].

For more information on Ultra Benfotiamine, visit douglaslabs.com

† These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

Manufactured by Douglas Laboratories 600 Boyce Road Pittsburgh, PA 15205 800-245-4440 douglaslabs.com



3

You trust Douglas Laboratories.
Your patients trust you.

© 2013 Douglas Laboratories. All Rights Reserved